

NewSil®

Imparts 100% plant available silicon



nutritional range

Group 2 Fertiliser

Reg. No. B5030 Act No. 36 of 1947

APPLICATION		
Crop	Application Rate	General
Nursery seedlings	50 ml/100 ℓ	Apply fortnightly
Vegetable crops	200 ml/100 ℓ	After planting; At fruit set Fortnightly thereafter
Tree crops	200 ml/100 ℓ	Before flowering; Full bloom, fruit set; Every 2-3 weeks thereafter
Row crops	200-400 ml/100 ℓ	After emergence; 5 leaf stage 2-4 weeks later; 6-8 weeks later

INSTRUCTIONS FOR USE

1. Maintain agitation while filling up.
2. Fertilisers and pesticides should be added after **NewSil®**.
3. For maximum uptake, spray in slow drying conditions and not to the point of run-off.
4. Repeat the foliar spray 3-4 times with of 14-21 day intervals.
5. **NB: Dilute as accurate as possible. Don't over dose!**

Conditions of Sale

Every care is taken during the manufacture of this product. As no control can be exercised over storage, handling, mixing, applications, use, weather, plant or soil conditions, no responsibility or liability for any failure in performance, losses, damage or injuries will be accepted under any circumstances whatsoever.

Distributed by:
realIPM South Africa (Pty) Ltd
Co. Reg. 2012/034646/07
73 Highlands Rd, Grabouw, 7160, RSA
info@realipm.co.za
www.realipm.co.za

Registered by:
Juniflor Finance & Control (Pty) Ltd
Co. Reg. 1998/024148/07
P.O. Box 2462, Wilropark,
Krugersdorp, RSA, 1731

Ortho silicic acid based fertiliser for the increase in resistance to biotic and abiotic plant stresses, nutrient uptake and crop quality

NUTRIENT ANALYSIS			
Macro Elements – g/ℓ			
Si	K	B	Cl
6.00	6.78	2.00	21.13
SG: 1.11		pH: 2.0	

IMPORTANT NOTE

Add **NewSil®** to water and **NOT** the other way around!



5 ℓ 25 ℓ 230 ℓ

SHAKE WELL BEFORE USE

Batch No.